

APA242Ra01 100µg

Active Nesfatin 1 (NES1)

Organism Species: *Rattus norvegicus (Rat)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Pro26~Leu106

Tags: Two N-terminal Tags, His-tag and GST-tag

Purity: >98%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 39.5kDa

Accurate Molecular Mass: 39kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

PIDVD KTKVHNVEPV ESARIEPPDT
GLYYDEYLKQ VIEVLETDPH FREKLQKADI EEIRSGRLSQ ELDLVSHKVR
TRLDEL

[ACTIVITY]

Nucb2 (Nucleobindin-2), a calcium-binding protein, is further cleaved into NES1 (Nesfatin-1). NES1 is an anorexigenic peptide and seems to participate in hypothalamic pathways regulating food intake and energy homeostasis, acting in a leptin-independent manner. GADD45A (Growth arrest and DNA damage-inducible protein GADD45 alpha) has been identified as an interactor of Nucb2. Besides, Growth Arrest And DNA Damage Inducible Protein Beta (GADD45b) has been identified as an interactor of NES1, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat NES1 and recombinant rat GADD45b. Briefly, NES1 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100µL were then transferred to GADD45b-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-NES1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of NES1 and GADD45b was shown in Figure 1, and this effect was in a dose dependent manner.




Figure 1. The binding activity of NES1 with GADD45b.

[IDENTIFICATION]




Figure 2. SDS-PAGE

Sample: Active recombinant NES1, Rat




Figure 3. Western Blot

Sample: Recombinant NES1, Rat;

Antibody: Rabbit Anti-Rat NES1 Ab (PAA242Ra01)

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.