

PAA372Ra01

Polyclonal Antibody to Gelsolin (GS)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ PRODUCT INFORMATION ]

**Immunogen:** GS, Rat **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

**Immunoglobulin Type**: IgG **UOM**: 100μg

## [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant GS (Ala432~Ala780) expressed in *E.coli*.

Accession No.: RPA372Ra01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSG S-AAQHGMDDD GTGQKQIWRI EGSNKVLVDP ATYGQFYGGD SYIILYNYRH GGRQGQIIYN WQGAQSTQDE VAASAILTAQ LDEELGGTPV QSRVVQGKEP AHLMSLFGGK PMIIYKGGTS RDGGQTTPAS TRLFQVRASS SGATRAVEVM PKAGALNSND AFVLKTPSAA YLWVGTGASD AEKTGALELL KVLRAQHVQV EEGSEPDGFW EALGGKTAYR TSPRLKDKKM DAHPPRLFAC SNRIGRFVIE EVPGELMQED LATDDVMLLD TWDQVFVWVG KDSQEEEKTE ALTSAKRYIE TDPANRDRRT PITVVRQGFE PPSFVGWFLG WDDDYWSVDP I DRAI AFI AA

#### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against GS. It has been selected for its ability to recognize GS in immunohistochemical staining and western blotting.

#### [APPLICATIONS]

Western blotting: 1:100-400




Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [ IMAGES ]



Used in DAB staining on fromalin fixed paraffin- embedded kidney tissue



Used in Western Blot ,Sample:

Recombinant GS,Rat

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant GS (Ala432~Ala780) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer



for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.