

**PAA099Rb52**

**Polyclonal Antibody to Matrix Metalloproteinase 13 (MMP13)**

**Organism Species: *Oryctolagus cuniculus* (Rabbit)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## [ **PRODUCT INFORMATION** ]

**Immunogen:** MMP13, Rabbit

**Clonality:** Polyclonal

**Host:** Cavia


**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg



*Sample: Recombinant MMP13, Rabbit*

## [ **IMMUNOGEN INFORMATION** ]

**Immunogen:** Recombinant MMP13 (Ser287~Gln449) expressed in *E. coli*.

**Accession No.:** RPA099Rb02

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSGS- SLSL DAITSLRGET MIFKDRFFWR LHPQQVDAEL FLTKSFWPEL  
PNRIDAAAYEH PARDLIFIFR GKKFWAPNGY DILEGYPQKL SELGFPREVK KISAAVHFED  
TGKTLFFSGN QVWSYDDTNH TMDQDYPRLI EEEFPGIGGK VDAVYEKNGY IYFFNGPIQ

## **[ ANTIBODY SPECIFICITY ]**

The antibody is a cavia polyclonal antibody raised against MMP13. It has been selected for its ability to recognize MMP13 in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## **[ CONTENTS ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## **[ QUALITY CONTROL ]**

**Content:** The quality control contains recombinant MMP13 (Ser287~Gln449) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.  
5uL per well when used in enhanced chemiluminescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

## **[ STORAGE ]**

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.