#### RPD195Mu01 50µg Recombinant Collagen Type III Alpha 1 (COL3a1) **Organism Species: Mus musculus (Mouse)** Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

kDa 70

44

33 26

22

18

14

10

#### [PROPERTIES]

Residues: Asp1224~Lys1371 Tags: N-terminal His-Tag Accession: Q7TT32 Host: E. coli Subcellular Location: Secreted, extracellular space, extracellular matrix. **Purity: >95%** Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). Formulation: Supplied as lyophilized form in 20mM Tris, 15% SDS-PAGE 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. Predicted isoelectric point: 6.4 Predicted Molecular Mass: 18.3kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

## [USAGE]

Reconstitute in ddH<sub>2</sub>O.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

# [<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

DFKINTE EIMSSLKSVN GQIESLISPD GSRKNPARNC RDLKFCHPEL KSGEYWVDPN QGCKMDAIKV FCNMETGETC INASPMTVPR KHWWTDSGAE KKHVWFGESM NGGFQFSYGP PDLPEDVVDV QLAFLRLLSS RASQNITYHC K

## [REFERENCES]

- 1. Christner P.J., et al. (2003) Biochem. Biophys. Res. Commun. 303:406-412.
- 2. Spiess K., et al. (2007) Connect. Tissue Res. 48:99-108.
- 3. Goldberg S.R., et al. (2007) J. Surg. Res. 143:27-34.
- 4. Carter R., et al. (2009) J. Surg. Res. 156:90-94.