

P93297Hu01 Cytochrome P450 1B1 (CYP1B1) **Organism: Homo sapiens (Human)** Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES

3th Edition (Revised in February, 2012)

| Human CYP1B1 | <u>kDa</u> | [DESCRIPTION] |
|--------------|------------|--|
| - | 94 | Protein Names: Cytochrome P450 1B1 |
| | 66.2 | Gene Names: CYP1B1 |
| | | Size: 100µg |
| *** | 45 | Source: Recombinant |
| | | Expression Host: E.coli |
| | | Function: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver |
| == | 33 | microsomes, this enzyme is involved in an NADPH-dependent electron transport |
| | | pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, |
| - | 26 | fatty acids, and xenobiotics. |
| | | Subcellular Location: Endoplasmic reticulum membrane; Peripheral membrane protein. |
| | | Microsome membrane; Peripheral membrane protein. |
| | 20 | Tissue Specificity: Expressed in many tissues. |
| | | [PROPERTIES] |
| | | Residues: Asp374~Phe516 (Accession # Q16678), with a N-terminal His-tag. |
| | 14 4 | Grade & Purity: >97%, 17.9 kDa as determined by SDS-PAGE reducing conditions. |

15% Tris-glycine SDS-PAGE **Endotoxin Level:** <1.0 EU per 1µg (determined by the LAL method).

Form & Buffer: Supplied as lyophilized form in PBS, pH 7.4.





Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 17.9 kDa

[PREPARATION]

Reconstitute in PBS.

[STORAGE AND STABILITY]

Storage: Store at 4°C for short time storage (1-2 weeks). Aliquot and store at -20°C or -80°C for long term storage.

Avoid repeated freeze/thaw cycles.

Valid period: 12 months stored at -80°C.

[BACKGROUND]

The target protein is fused with a His-tag and its sequence is listed below. The first Met is an initiator amino acid. Moreover, Gly and Ser are added to improve the flexibility of N-terminus at both ends of the His-tag, which will increase the chelating ability of the tag to Ni-Sepharose during purification.

MGHHHHHHSGSEF-DQPNLPY VLAFLYEAMR FSSFVPVTIP HATTANTSVL GYHIPKDTVV FVNQWSVNHD PLKWPNPENF DPARFLDKDG LINKDLTSRV MIFSVGKRRC IGEELSKMQL FLFISILAHQ CDFRANPNEP AKMNFSYGLT IKPKSF

