Coud-Clone Corp.

RPA234Hu01 10µg Recombinant Immunoglobulin G4 (IgG4) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Gln222~Lys327	kDa		
Tags: Two N-terminal Tags, His-tag and T7-tag	94 66.2		
Accession: P01861		1	
Host: E. coli	45		
Subcellular Location: Secreted.	33	-	
Purity: >95%	26	-	
Endotoxin Level: <1.0EU per 1µg			
(determined by the LAL method).	20		
Formulation: Supplied as lyophilized form in 100mM		-	
NaHCO3, 500mM NaCl, pH8.3, containing 1mM EDTA,	14.4		
1mM DTT, 0.01% sarcosyl, 5% trehalose, and			
preservative.	15% SDS-PAGE		
Predicted isoelectric point: 6.5			
Predicted Molecular Mass: 15.9kDa			
Applications: SDS-PAGE; WB; ELISA; IP.			
(May be suitable for use in other assays to be determined by the end user.)			

[<u>USAGE</u>]

Reconstitute in ddH₂O.

Cloud-Clone Corp.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

QPREPQVYT LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTPPVLDS DGSFFLYSRL TVDKSRWQEG NVFSCSVMHE ALHNHYTQKS LSLSLGK

[REFERENCES]

- 1. Ellison J.W., et al. (1981) DNA 1:11-18.
- 2. Pink J.R.L., et al. (1970) Biochem. J. 117:33-47.
- 3. Chen R., et al. (2009) J. Proteome Res. 8:651-661.
- 4. Davies A.M., et al. (2013) Mol. Immunol. 54:1-7.