

**MAA644Ra22** 

Monoclonal Antibody to CCAAT/Enhancer Binding Protein Alpha (CEBPa)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: N/A

Purification: Protein A + Protein G affinity chromatography

Clone number: C2

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 100µl

Cross Reactivity: Human

Applications: WB; IHC; ICC; IP.

## [ IMMUNOGEN ]

Immunogen: Recombinant CEBPa (Ala72~Ala358) expressed in E.coli

Accession No.: RPA644Ra01

## [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

## [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined



by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]



Western Blot; Sample: Lane1: U937 cell lysate; Lane2: THP1 cell lysate Primary Ab: 0.2µg/ml Mouse Anti-Rat CEBPa Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Mouse IgG

Polyclonal Antibody (Catalog: SAA544Mu19)

Western Blot; Sample: Lane1: U937 cell lysate; Lane2: THP1 cell lysate Primary Ab: 0.2µg/ml Mouse Anti-Rat CEBPa Antibody Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.