

**MAA480Po21** 

**Monoclonal Antibody to Myoglobin (MYO)** 

Organism Species: Sus scrofa; Porcine (Pig)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



# [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C1

Traits: Liquid

Concentration: 0.85mg/ml

**UOM:** 100µl

Cross Reactivity: Mouse; Rat

Applications: WB; IHC; ICC; IP.

## [ IMMUNOGEN ]

Immunogen: Recombinant MYO (Met1~Gly154) expressed in E.coli

Accession No.: RPA480Po01

#### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined


# Coud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ IDENTIFICATION ]



Western Blot; Sample: Lane1: Porcine
Heart lysate; Lane2: Rat Heart lysate;
Lane3: Mouse Heart lysate Primary Ab:
0.15µg/ml Mouse Anti-Porcine MYO
Antibody Second Ab: 0.2µg/mL HRPLinked Caprine Anti-Mouse IgG
Polyclonal Antibody (Catalog:
SAA544Mu19)



DAB staining on IHC-P; Sample:
Porcine Cardiac Muscle Tissue;
Primary Ab: 20µg/ml Mouse AntiPorcine MYO Antibody Second Ab:
2µg/mL HRP-Linked Caprine AntiMouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)

DAB staining on IHC-P;


Sample: Porcine Skeletal muscle
Tissue;

Primary Ab: 20µg/ml Mouse AntiPorcine MYO Antibody


Second Ab: 2µg/mL HRP-Linked

Caprine Anti-Mouse IgG Polyclonal
Antibody

(Catalog: SAA544Mu19)



DAB staining on IHC-P;
Sample: Porcine Cardiac Muscle
Tissue
Primary Ab: 20µg/ml Mouse AntiPorcine MYO Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked



DAB staining on IHC-P;
Sample: Porcine Skeletal muscle
Tissue
Primary Ab: 20µg/ml Mouse AntiPorcine MYO Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked



Caprine Anti-Mouse IgG Polyclonal

Caprine Anti-Mouse IgG Polyclonal

Antibody

Antibody

(Catalog: SAA544Mu19)

(Catalog: SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.