

PAL260Mu01 Polyclonal Antibody to Brother Of CDO Homolog (BOC) Organism Species: *Mus musculus (Mouse) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

### [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

**UOM:** 100µl

Cross Reactivity: N/A

Applications: WB; IHC; ICC; IP.

#### [ IMMUNOGEN ]

Immunogen: Recombinant BOC (Lys668~Thr866) expressed in E.coli

Accession No.: RPL260Mu01

#### [APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

## Cond-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [IDENTIFICATION]





DAB staining on IHC-P; DAB staining on IHC-P; Samples: Mouse Pancreas Tissue; Samples: Mouse Spleen Tissue; Primary Ab: 10µg/ml Rabbit Anti-Mouse Primary Ab: 10µg/ml Rabbit Anti-Mouse BOC Antibody BOC Antibody Second Ab: 2µg/mL HRP-Linked Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Caprine Anti-Rabbit IgG Polyclonal Antibody Antibody (Catalog: SAA544Rb19) (Catalog: SAA544Rb19)

## [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.