

PAC413Mu01

Polyclonal Antibody to Calcitonin Receptor (CTR) Organism Species: *Mus musculus (Mouse)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

Cond-Clone Corp.

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.66mg/ml

UOM: 100µl

Cross Reactivity: Rat

Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant CTR (Ala42~Leu170) expressed in E.coli

Accession No.: RPC413Mu01

[APPLICATIONS]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

Contend Cloud - Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]






Figure. Western Blot; Sample:

Recombinant CTR, Mouse.



lysate;

CTR Antibody



DAB staining on IHC-P;

Samples: Mouse Kidney Tissue; Western Blot; Sample: Mouse Kidney Primary Ab: 20?g/ml Rabbit Anti-Mouse **CTR** Antibody Primary Ab: 1µg/ml Rabbit Anti-Mouse Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Second Ab: 0.2µg/mL HRP-Linked Antibody Caprine Anti-Rabbit IgG Polyclonal (Catalog: SAA544Rb19)

Antibody (Catalog: SAA544Rb19)

[<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.