

PAA853Hu01

Polyclonal Antibody to Caspase 8 (CASP8)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[**PROPERTIES**]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µL

Cross Reactivity: Mouse;Rat;Porcine

Applications: WB; IHC; ICC/IF

[**IMMUNOGEN**]

Immunogen: Recombinant CASP8 (Ser217~Asp384) expressed in *E.coli*

Accession No.: RPA853Hu01

[**APPLICATIONS**]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-30µg/mL;

Immunofluorescence:5-20µg/mL;

Optimal working dilutions must be determined by end user.

[**FORMULATION**]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


[IDENTIFICATION]




DAB staining on IHC-P; Samples: Human Kidney Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human CASP8 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Sample: Rat Small intestine Tissue Primary Ab: 10µg/ml Rabbit Anti-Human CASP8 Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Samples: Human Liver cancer Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human CASP8 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P; Sample: Human Lymphoma Tissue Primary Ab: 20µg/ml Rabbit Anti-Human CASP8 Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked




DAB staining on IHC-P; Samples: Human Colon Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human CASP8 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Sample: Human Amygdalitis Tissue Primary Ab: 20µg/ml Rabbit Anti-Human CASP8 Antibody Control: Used PBS instead of primary antibody Second Ab: 2?g/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)




Western Blot; Sample: Lane1: HepG2 cell lysate; Lane2: Hela cell lysate; Lane3: Raji cell lysate; Lane4: Porcine Spleen lysate; Lane5: Mouse Lung lysate; Lane6: Rat Spleen lysate
Primary Ab: 0.2 μ g/ml Rabbit Anti-Human CASP8 Antibody
Second Ab: 0.2 μ g/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb19)




FITC staining on IF;
Samples: Human HepG2 cell;
Primary Ab: 20 μ g/ml Rabbit Anti-Human CASP8 Antibody
Second Ab: 1.5 μ g/ml FITC-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)



Knockout Varification:
Lane 1: Wild-type Hela cell lysate; Lane 2: CASP8 knockout Hela cell lysate;
Predicted MW: 26-32,46,54-62kd
Observed MW: 35kd
Primary Ab: 1 μ g/ml Rabbit Anti-Human CASP8 Antibody
Second Ab: 0.2 μ g/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb19) Selected



FITC staining on IF;
Sample: HepG2 cell
Primary Ab: 20 μ g/ml Rabbit Anti-Human CASP8 Antibody
Second Ab: 2 μ g/ml FITC-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb11)

(Catalog: SAA544Rb18)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.