

PAA308Mu01

Polyclonal Antibody to Galectin 8 (GAL8)
Organism Species: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [ PRODUCT INFORMATION ]

**Immunogen:** GAL8, Mouse **Purification:** Affinity Chromatography.

Clonality: Polyclonal Applications: WB, ICC, IHC-P, IHC-F, ELISA

Host: Rabbit Concentration: 200µg/mL

**Immunoglobulin Type**: IgG **UOM**: 100μg

### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant GAL8 (Met1~Trp316) expressed in E.coli.

Accession No.: RPA308Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGS- MLSLNNLQNI IYNPIIPYVG TITEQLKPGS LIVIRGHVPK DSERFQVDFQ LGNSLKPRAD VAFHFNPRFK RSSCIVCNTL TQEKWGWEEI TYDMPFRKEK SFEIVFMVLK NKFQVAVNGR HVLLYAHRIS PEQIDTVGIY GKVNIHSIGF RFSSDLQSME TSALGLTQIN RENIQKPGKL QLSLPFEARL NASMGPGRTV VIKGEVNTNA RSFNVDLVAG KTRDIALHLN PRLNVKAFVR NSFLQDAWGE EERNITCFPF SSGMYFEMII YCDVREFKVA INGVHSLEYK HRFKDLSSID TLSVDGDIRL LDVRSW

### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against GAL8. It has been selected for its ability to recognize GAL8 in immunohistochemical staining and western blotting.

### [APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500




Immunohistochemistry in formalin fixed frozen section: 1:50-500


Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

# [IMAGES]



Used in DAB staining on fromalin fixed paraffin-embedded testis tissue



Used in Western Blot, Sample:
Recombinant GAL8, Mouse

# [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant GAL8 (Met1~Trp316) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN<sub>3</sub> 0.02%.

### [STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.