

PAB655Mu02

Polyclonal Antibody to Interleukin-18 Receptor Beta (IL18R Beta)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µL

Cross Reactivity: Rat

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant IL18R Beta (Gln117~Arg240 (Accession # Q9Z2B1)) expressed in

E.coli

Accession No.: RPB655Mu02

#### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined



by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]



DAB staining on IHC-P; Samples: DAB staining on IHC-P;

Mouse Kidney Tissue; Primary Ab: Samples: Mouse Colon Tissue;

20?g/ml Rabbit Anti-Mouse IL18RAP Primary Ab: 20?g/ml Rabbit Anti-Mouse

Antibody Second Ab: 2µg/mL HRP- IL18RAP Antibody

Linked Caprine Anti-Rabbit IgG Second Ab: 2µg/mL HRP-Linked

Polyclonal Antibody (Catalog: Caprine Anti-Rabbit IgG Polyclonal

SAA544Rb19) Antibody

(Catalog: SAA544Rb19)

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.