



pA91868Rb51 Polyclonal Antibody to Metallothionein 2 (MT2) Organism: Oryctolagus cuniculus (Rabbit) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

7th Edition (Revised in May, 2013)

Western Blot

kDa

[PRODUCT INFORMATION]

Immunogen: MT2, Rabbit

Clonality: Polyclonal

Host: Cavia

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100μg Sample: Recombinant MT2, Rabbit

75 50 37 25 20 15 10 ←8kDa

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant MT2 (Met1~Ala62) expressed in E.coli.

USCN Accession No.: rP91868Rb01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSG SEF- MDPNCSCAAA GDSCTCANSC TCKACKCTSC KKSCCSCCPP

GCAKCAQGCI CKGASDKCSC CA





[ANTIBODY SPECIFITY]

The antibody is a cavia polyclonal antibody raised against MT2. It has been selected for its ability to recognize MT2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant MT2 (Met1~Ala62) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.