

**PAC919Hu01**

**Polyclonal Antibody to Platelet Derived Growth Factor D (PDGFD)**

**Organism Species: Homo sapiens (Human)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

9th Edition (Revised in Jul, 2013)

## **[ PRODUCT INFORMATION ]**

**Immunogen:** PDGFD, Human

**Clonality:** Polyclonal

**Host:** Rabbit

**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg

## **[ IMMUNOGEN INFORMATION ]**

**Immunogen:** Recombinant PDGFD (Arg19~Arg370) expressed in *E.coli*.

**Accession No.:** RPC919Hu01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF- RD TSATPQSASI  
KALRNANLRR DESNHLTDLY RRDETIQVKG NGYVQSPRFP NSYPRNLLLT WRLHSQENTR  
IQLVFDNQFG LEEAENDICR YDFVEVEDIS ETSTIIRGRW CGHKEVPPRI KSRTNQIKIT  
FKSDDYFVAK PGFKIYYSLL EDFQAAAASE TNWESVTSSI SGVSYNSPSV TDPTLIADAL  
DKKIAEFDTV EDLLKYFNPE SWQEDLENMY LDTPRYRGRS YHDRKSKVDL DRLNDDAKRY  
SCTPRNYSVN IREELKLANV VFFPRCLLVQ RCGNCGCGT VNWRSTCNS GKTVKKYHEV  
LQFEPGHIKR RGRAKTMALV DIQLDHHERC DCICSSRPPR

## **[ ANTIBODY SPECIFICITY ]**

The antibody is a rabbit polyclonal antibody raised against PDGFD. It has been selected for its ability to recognize PDGFD in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500


Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100


Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## [ **IMAGES** ]



Used in DAB staining on formalin fixed  
paraffin-embedded ovary tissue



Used in Western Blot, Sample:  
Recombinant PDGFD, Human

## [ **CONTENTS** ]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02%  $\text{NaN}_3$ , 50% glycerol.

## [ **QUALITY CONTROL** ]

**Content:** The quality control contains recombinant PDGFD (Arg19~Arg370) disposed in loading buffer.

**Usage:** 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.  
5uL per well when used in enhanced chemiluminescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

**Loading Buffer:** 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%,  $\text{NaN}_3$  0.02%.

## [ **STORAGE** ]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.