

PAD357Hu01

Polyclonal Antibody to Protein Kinase, AMP Activated Alpha 2 (PRKAa2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 1mg/ml

**UOM**: 100µg(100µl)

Cross Reactivity: Mouse; Rat; Hamster; Canine; Porcine; Bovine; Caprine; Ovine; Gallus

Applications: WB; ICC; IHC-P; IHC-F; ELISA.

#### [ IMMUNOGEN ]

Immunogen: Recombinant PRKAa2 (Asp252~Ala493) expressed in E.coli

Accession No.: RPD357Hu01

## [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no



obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]





Figure. Western Blot; Sample: Recombinant PRKAa2, Human.



DAB staining on IHC-P; Samples: Human Kidney Tissue

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.