## ے <u>Cloud-Clone Co</u>rp.

PAA198Hu01 Polyclonal Antibody to Ribonuclease P (RNASEP) Organism Species: Homo sapiens (Human) *Instruction manual* 

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

### [PRODUCT INFORMATION]

Immunogen: RNASEP, Human Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

#### [ IMMUNOGEN INFORMATION ]

Immunogen: Recombinant RNASEP (Cys49~Asn258) expressed in E.coli.

Accession No.: RPA198Hu01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGS- CG ILSEELKNLV MNTGPYYFVK NLPLHELITP EFISTFIKKG SCYALTYNTH IDEDNTVALL PNGKLILSLD KDTYEETGLQ GHPSQFSGRK IMKFIVSIDL MELSLNLDSK KYERISWSFK EKKPLKFDFL LAWHKTGSEE STMMSYFSKY QIQEHQPKVA LSTLRDLQCP VLQSSELEGT PEVSCRALEL FDWLGAVFSN VDLNNEPN

## [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against RNASEP. It has been selected for its ability to recognize RNASEP in immunohistochemical staining and western blotting.


## [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500

# Cloud-Clone Corp.


Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

### [<u>IMAGES</u>]



Used in DAB staining on fromalin fixed paraffin-embedded Brain tissue

### [CONTENTS]



Used in Western Blot, Sample: Recombinant RNASEP, Human

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant RNASEP (Cys49~Asn258) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

## [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.