

CSI245Rb01 Primary Rabbit Spermatogonium Cells (SPC) Organism Species: Oryctolagus cuniculus (Rabbit) Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Revised in Aug, 2023)

### [ DESCRIPTION ]

Cell Type: Spermatogonium cell Synonyms: SPC Species: Oryctolagus cuniculus (Rabbit) Tissue Source: Testis Size: >5×10<sup>5</sup>cell/vial

### [PROPERTIES]

Cell activity: >85% (Viability by Trypan Blue Exclusion).

Formulation: Frozen 1 mL or T25 flask.

Biosafety: Negative for HIV-1, HBV, HCV, mycoplasma, bacteria, yeast and fungi.

**Applications:** For research use only. It is not approved for human or animal use, or for application in clinical diagnostic procedures.

Growth Properties: Adherent

# [ CONTENTS ]

Form & Buffer: Supplied as solution form in frozen stock solution, containing 90% FBS+10% DMSO.

## [USAGE]

Upon receiving the cells in a T-25 flask at room temperature, immediately transfer the cells to 37°C, 5% incubator; the cells in vials, directly and immediately transfer the cells from dry ice to liquid nitrogen.

#### **Culture conditions:**

 $\mathsf{DMEM+5\%FBS+1\%Spermatogonium\ Cell\ Growth\ Supplement+1\%Penicillin-Streptomycin\ Solution}$ 

Temperature: 37°C

Condition: 95% air, 5% carbon dioxide

#### Cell recovery:

After receiving the cells, shake at 37°C in a water bath until completely dissolved, transfer to a 15 ml centrifuge tube, add 3-5 times complete culture solution, 1000 rpm for 5 min, discard the supernatant, and place in a T25 flask for culture.

#### Cell passage:

- 1. Cell passage when cell growth at 85-95%.
- 2. Discard the medium and wash with PBS 1-2 times.

# Cloud-Clone Corp.

- 3. Add 1 ml of Trypsin at 37°C, observe the cell under the microscope. If the cells are retracted and rounded, pat the culture flask to let the cells fall off. Stop digestion by adding 2 ml of complete medium containing 10% serum. Make it a single cell suspension.
- 4. Add the fresh medium to resuspend the cells. Unless otherwise stated, the recommended ratio of primary cells is 1/2.

# [STORAGE]

Freeze of the liquid nitrogen (90% FBS +10% DMSO).

# [Shipping]

Dry ice.

# [IMPORTANT NOTE]

The cell is for research use only, and we will not be responsible for any issue if the cell was used in clinical diagnostic or any other procedures.

## [Figure]




Figure 1

Figure 2

Figure 1 Morphology of Rabbit Spermatogonium Cells (Optical microscope,×100) Figure 2 Morphology of Rabbit Spermatogonium Cells (Optical microscope,×200)