

RPJ640Hu01 50µg Recombinant Anillin (ANLN) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Cond-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Cys712~Pro1124

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose .

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.5

Predicted Molecular Mass: 50.1kDa

Accurate Molecular Mass: 50&35kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in ddH_2O to a concentration of 0.1-0.2 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

Contend Clone Corp.

CQVNIKQKM QELNNEINMQ QTVIYQASQA LNCCVDEEHG KGSLEEAEAE RLLLIATGKR TLLIDELNKL KNEGPQRKNK ASPQSEFMPS KGSVTLSEIR LPLKADFVCS TVQKPDAANY YYLIILKAGA ENMVATPLAS TSNSLNGDAL TFTTTFTLQD VSNDFEINIE VYSLVQKKDP SGLDKKKKTS KSKAITPKRL LTSITTKSNI HSSVMASPGG LSAVRTSNFA LVGSYTLSLS SVGNTKFVLD KVPFLSSLEG HIYLKIKCQV NSSVEERGFL TIFEDVSGFG AWHRRWCVLS GNCISYWTYP DDEKRKNPIG RINLANCTSR QIEPANREFC ARRNTFELIT VRPQREDDRE TLVSQCRDTL CVTKNWLSAD TKEERDLWMQ KLNQVLVDIR LWQPDACYKP IGKP

Figure . Gene Sequencing (extract)

Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.