

RPA014Hu02 10µg

Recombinant Bone Morphogenetic Protein 4 (BMP4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Ser22~Arg408

Tags: N-terminal His Tag

Subcellular Location: Secreted

**Purity:** > 97%

Traits: Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.0

Predicted Molecular Mass: 48.2kDa

**Accurate Molecular Mass:** 48kDa as determined by SDS-PAGE reducing conditions.

### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

#### [ SEQUENCE ]



		SHASLIPET	GKKKVAEIQG	HAGGRRSGQS
HELLRDFEAT	LLQMFGLRRR	PQPSKSAVIP	DYMRDLYRLQ	SGEEEEEQIH
STGLEYPERP	ASRANTVRSF	HHEEHLENIP	GTSENSAFRF	LFNLSSIPEN
EVISSAELRL	FREQVDQGPD	WERGFHRINI	YEVMKPPAEV	VPGHLITRLL
DTRLVHHNVT	RWETFDVSPA	VLRWTREKQP	NYGLAIEVTH	LHQTRTHQGQ
HVRISRSLPQ	GSGNWAQLRP	LLVTFGHDGR	GHALTRRRRA	KRSPKHHSQR
ARKKNKNCRR	HSLYVDFSDV	GWNDWIVAPP	GYQAFYCHGD	CPFPLADHLN
STNHAIVQTL	VNSVNSSIPK	ACCVPTELSA	ISMLYLDEYD	KVVLKNYQEM
VVEGCGCR				

# [ IDENTIFICATION ]



# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.