

RPB134Ra01 50ug Recombinant Chondroitin Sulfate Proteoglycan 4 (CSPG4) Organism Species: *Rattus norvegicus (Rat) Instruction manual* 

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

# Cond-Clone Corp.

# [PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Asn1135~Gln1444

Tags: N-terminal His Tag

Subcellular Location: Membrane, Extracellular matrix

**Purity:** > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.1% SKL, 1mM DTT, 5% Trehalose and

Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 34.6kDa

Accurate Molecular Mass: 35kDa as determined by SDS-PAGE reducing conditions.

#### [<u>USAGE</u>]

Reconstitute in PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ <u>SEQUENCE</u> ]

# Cond-Clone Corp.

NSSSLV VPQGQQGTID TAVLHLDTNL DIRSGNEVHY HVTAGPHWGQ LLRDGQSVTS FSQRDLLDGA ILYSHNGSLS PQDTLALSVA AGPVHTSTVL QVTIALEGPL APLQLVQHKR IYVFQGEAAE IRRDQLEVVQ EAVLPADIMF SLRSPPNAGY LVMVSHGASA DGPPSLDPVQ RFSQEAINSG RVLYLHSRPG AWSDSFSLDV ASGLGDPLEG ISVELEVLPT VIPLDVQNFS VPEGGTRTLA PPLIQITGPY LGTLPGLVLQ VLEPPQHGAL QKEDRPODGT LSTFSWREVE EQLIRYVHDG SETQ

## [IDENTIFICATION]




Figure. SDS-PAGE

#### [IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.