RPC366Hu01 50µq **Recombinant Cytidine Deaminase (CDA)** Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

C Cloud-Clone Corp.

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Ala2~Gln146 **Tags:** Two N-terminal Tags, His-tag and T7-tag Accession: P32320 Host: E. coli **Purity:** >95% Endotoxin Level: <1.0EU per 1µg (determined by the LAL method). **Formulation:** Supplied as lyophilized form in 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. 15% SDS-PAGE Predicted isoelectric point: 6.8 Predicted Molecular Mass: 19.9kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

[USAGE]

Reconstitute in ddH₂O.



Coud-Clone Corp.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

AQKRPACTL KPECVQQLLV CSQEAKKSAY CPYSHFPVGA ALLTQEGRIF KGCNIENACY PLGICAERTA IQKAVSEGYK DFRAIAIASD MQDDFISPCG ACRQVMREFG TNWPVYMTKP DGTYIVMTVQ ELLPSSFGPE DLQKTQ

[REFERENCES]

- 1. Laliberte J., Momparler R.L. (1994) Cancer Res. 54:5401-5407.
- 2. Demontis S., et al. (1998) Biochim. Biophys. Acta 1443:323-333.
- 3. Gran C., et al. (1998) Blood 91:4127-4135.
- 4. Kuhn K., et al. (1993) Biochem. Biophys. Res. Commun. 190:1-7.