

RPB215Hu01 100µg

Recombinant Fibrinogen Beta (FGb)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[**PROPERTIES**]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Gly45~Gln491

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.6

Predicted Molecular Mass: 52.0kDa

Accurate Molecular Mass: 52kDa as determined by SDS-PAGE reducing conditions.

[**USAGE**]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[**STORAGE AND STABILITY**]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.


Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

KKREEAPSLR PAPPPIISGGG YRARPAKAAA TQKKVERKAP DAGGCLHADP
 DLGVLCPGTC QLQEALLQQE RPIRNSVDEL NNNVEAVSQT SSSSFQMYL
 LKDLWQKRQK QVKDNENVVN EYSSELEKHQ LYIDETVNSN IPTNLRVLRS
 ILENLRSKIQ KLESDVSAQM EYC RTPCTVS CNIPVVSQKE CEEIIRKGGG
 TSEMYLIQPD SSVKPYRVYC DMNTENGGWT VIQNRQDGSV DFGRKWDPYK
 QGFGNVATNT DGKNYCGLPG EYWLGNDKIS QLTRMGPTL LIEMEDWKGD
 KVKAHYGGFT VQNEANKYQI SVNKYRGTAG NALMDGASQL MGENRTMTIH
 NGMFFSTYDR DNDGWLTS DP RKQCSKEDGG GWWYNRCHAA NPNGRYYWGG
 QYTWDMAKHG TDDGVVWMNW KGSWYSMRKM SMKIRPFFPQ Q

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.