

RPE540Ra01 1 Recombinant Testicular Cell Adhesion Molecule 1 (TCAM1) Organism Species: *Rattus norvegicus (Rat) Instruction manual*

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression Host: *E.coli*

Residues: Ser226~Cys548

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.3

Predicted Molecular Mass: 39.3kDa

Accurate Molecular Mass: 39kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

Coud-Clone Corp.

SLLEV GMAETMSCEV VRVFPAQEAV

FRMFLEDQE LSPFSSWKGD AAWASATIQA METGDQELTC LVSVGPVEQK A RKPVHVYS FPPPVLEIED AYPQAGTDVN VTCSGHVLTS PSPTLRLQGS LN LSAPGEP AWLRFTAREE DDGRTLSCEA SLVVQGQRLV KTTKIQLHVL YKP RFQESD CPGNQIWVEG MDQMLACIPE GNPIPALVCI WNGMTFDLEV PQKA TQNHT GTYSCTATNS LGSVSKDIAV LVQGLHEGIS SSTIFIIIF TLGMA VITI ALYLNYQPCK RNGRKRTHRQ KEQNKGGERQ FSDIQAEECH AHLC

[IDENTIFICATION]

| 1 | kDa 70 |
|---|-----------|
| | 44 |
| - | 33 |
| | 26 |
| | 22 |
| | 18 |
| | 14 |
| | 10 |

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.