

RPA135Hu04 100µg Recombinant Thrombopoietin (TPO) Organism Species: *Homo sapiens (Human)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

Coud-Clone Corp.

[PROPERTIES]

Source: Prokaryotic expression Host: *E.coli* Residues: Ser22~Gly349 Tags: His and TrxA Taq Subcellular Location: Secreted Purity: > 80% Traits: Freeze-dried powder Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose . Original Concentration: 200µg/mL Applications: Positive Control; Immunogen; SDS-PAGE; WB. (May be suitable for use in other assays to be determined by the end user.) Predicted isoelectric point: 8.5

Predicted Molecular Mass: 55.4kDa

Accurate Molecular Mass: 55kDa as determined by SDS-PAGE reducing conditions.

[<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



| | | SPAPPACDL | RVLSKLLRDS | HVLHSRLSQC |
|------------|------------|------------|------------|------------|
| PEVHPLPTPV | LLPAVDFSLG | EWKTQMEETK | AQDILGAVTL | LLEGVMAARG |
| QLGPTCLSSL | LGQLSGQVRL | LLGALQSLLG | TQGRTTAHKD | PNAIFLSFQH |
| LLRGKVRFLM | LVGGSTLCVR | RAPPTTAVPS | RTSLVLTLNE | LPNRTSGLLE |
| TNFTASARTT | GSGLLKWQQG | FRAKIPGLLN | QTSRSLDQIP | GYLNRIHELL |
| NGTRGLFPGP | SRRTLGAPDI | SSGTSDTGSL | PPNLQPGYSP | SPTHPPTGQY |
| TLFPLPPTLP | TPVVQLHPLL | PDPSAPTPTP | TSPLLNTSYT | HSQNLSQEG |

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.