

RPB595Hu01 10ug

Recombinant Tyrosine Kinase 2 (Tyk2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



# [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Cys915~His1173

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Secreted, Cytoplasm

**Purity:** > 97%

Traits: Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.8

Predicted Molecular Mass: 33.5kDa

**Accurate Molecular Mass:** 34kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ SEQUENCE ]



CYDPTN DGTGEMVAVK ALKADCGPQH RSGWKQEIDI LRTLYHEHII KYKGCCEDQG EKSLQLVMEY VPLGSLRDYL PRHSIGLAQL LLFAQQICEG MAYLHAQHYI HRDLAARNVL LDNDRLVKIG DFGLAKAVPE GHEYYRVRED GDSPVFWYAP ECLKEYKFYY ASDVWSFGVT LYELLTHCDS SQSPPTKFLE LIGIAQGQMT VLRLTELLER GERLPRPDKC PCEVYHLMKN CWETEASFRP TFENLIPILK TVH

## [ IDENTIFICATION ]



### [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.