

APC374Mu61 100µg

Active Carboxylesterase 1 (CES1)

Organism Species: *Mus musculus (Mouse)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

[PROPERTIES]

Source: Eukaryotic expression.

Host: 293F cell

Residues: His19~Leu565

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 5% trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.9

Predicted Molecular Mass: 62.4kDa

Accurate Molecular Mass: 60kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.6) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were


observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```
HP SLPPVVHTVH GKVLGKYVTL EGFSQPVAVF
LGVPFAPKPL GSLRFAPPEP AEPWSFVKHT TSYPPLCYQN PEAAALRLAEL
FTNQRKIIPH KFSEDCLYLN IYTPADLTQN SRLPVMVWIH GGGLVIDGAS
TYDGVPLAVH ENVVVVVIQY RLGIWGFFST EDEHSRGNWG HLDQVAALHW
VQDNIANFGG NPGSVTIFGE SAGGESVSVL VLSPLAKNLF HRAIAQSSVI
FNPCLFGRAA RPLAKKIAAL AGCKTTTSAA MVHCLRQKTE DELLEVSLKM
KFGTVDFLGD PRESYPFLPT VIDGVLLPKA PEEILAEKSF NTPVYMVGIN
KHEFGWIIPM FLDPLSERK LDQKTAASIL WQAYPILNIS EKLIPAAIEK
YLGGTEDPAT MTDLFLDLIG DIMFGVPSVI VSRSHRDAGA PTYMYEYQYR
PSFVSDDRPQ ELLGDHADEL FSVWGAPFLK EGASEEEINL SKMVMKFWAN
FARNGNPNGE GLPHWPEYDQ KEGYLQIGVP AQAARHLKDK EVDFWTELRA
KETAERSSHR EHVEL
```

[**ACTIVITY**]

carboxylesterase 1(CES1) also known as Liver carboxylesterase 1 is a serine esterase and member of a large multigene carboxylesterase family. The protein Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl-CoA ester. Hydrolyzes the methyl ester group of cocaine to form benzoylecgonine. Thus, the recombinant mouse CES1 activity was measured by its ability to hydrolyze 4-Nitrophenyl acetate (4-NPA) to 4-Nitrophenol. The reaction was performed in 50 mM Tris, 150 mM NaCl, pH 7.5 (Assay Buffer), ainitiated by addition 50 μ L of various concentrations of CES1 (diluted by Assay Buffer) to 50 μ L of 2 mM Substrate 4-NPA (100 mM stock in Acetone, diluted by Assay Buffer). Incubated at 37 $^{\circ}$ C for 10min, then read at a wavelength of 400 nm.



4-Nitrophenol (product)mM/L	OD400nm
0.625	1.909
0.3125	1.032333333
0.15625	0.523333333
0.078125	0.269333333
0.0390625	0.144666667

Figure 1. The standard curve of 4-Nitrophenol

One unit of enzyme activity is defined as the 1µg of enzyme required to convert 1pmol of 4-Nitrophenyl acetate to 4-Nitrophenol in 1min at 37°C. The specific activity of recombinant mouse CES1 is 368.9 pmol/min/µg.

$$\text{Specific Activity (pmol/min/}\mu\text{g)} = \frac{\Delta OD * F}{T * N}$$

△OD=Adjusted for Substrate Blank

F=Conversion Factor(convert from standard curve of 4-Nitrophenol)

T= Time

N=Amount of enzyme

[IDENTIFICATION]




Figure 2. Gene Sequencing (extract)




Figure 3. SDS-PAGE

Sample: Active recombinant CES1, Mouse




Figure 4. Western Blot

Sample: Recombinant CES1, Mouse;

Antibody: Rabbit Anti-mouse CES1 Ab (PAC374Mu06)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.