

APF283Mu01 100µg
Active Carboxylesterase 3 (CES3)
Organism Species: *Mus musculus (Mouse)*
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Pro344~Leu565

Tags: N-terminal His-tag

Purity: >98%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.3

Predicted Molecular Mass: 28.7kDa

Accurate Molecular Mass: 29kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

PYIVGIN KQEFGWIIPT LMGYPLAEGK LDQKTANSLI WKSYPYTKIS ENMIPVVAEK YLGGTDDLTK KKDLEFDLMA
DVVFGVPSVI VSRSHRDAGA STYMYEFYR PSFVSAMRPK AVIGDHGDEI FSVFGSPFLK DGASEEETNL SKMVMKFWAN
FARNGNPNGG GLPHWPEYDQ KEGYLKIGAS TQAAQLKDK EVSFWAELRA KESAQRPSHR EHVEL

[ACTIVITY]

Carboxylesterase 3 (CES3) also known as Liver carboxylesterase 3 is a serine esterase and a member of a large family of carboxylesterases. The protein involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl-CoA ester. Hydrolyzes the methyl ester group of cocaine to form benzoylecgonine. Thus, the recombinant mouse CES3 activity was measured by its ability to hydrolyze 4-Nitrophenyl acetate (4-NPA) to 4-Nitrophenol. The reaction was performed in 50 mM Tris, pH 7.5 (Assay Buffer), initiated by addition 50 µL of various concentrations of CES3 (dilute by Assay Buffer) to 50 µL of 2 mM Substrate 4-NPA (100 mM stock in Acetone, dilute by deionized water). Incubated at 37 °C for 10min, then read at a wavelength of 400 nm




Figure 1. The standard curve of 4-Nitrophenol

One unit of enzyme activity is defined as the 1µg of enzyme required to convert 1pmol of 4-Nitrophenyl acetate to 4-Nitrophenol in 1min at 37°C. The specific activity of recombinant mouse CES3 is 900 pmol/min/µg.

$$\frac{\Delta OD * F}{T * N}$$

Specific Activity (pmol/min/µg) = $\frac{\Delta OD * F}{T * N}$

△OD=Adjusted for Substrate Blank

F=Conversion Factor(convert from standard curve of 4-Nitrophenol)

T= Time

N=Amount of enzyme

[IDENTIFICATION]




Figure 2. SDS-PAGE

Sample: Active recombinant CES3, Mouse




Figure 3. Western Blot

Sample: Recombinant CES3, Mouse;

Antibody: Rabbit Anti- Mouse CES3 Ab (PAF283Mu01)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.