

**APN576Mu61 50µg**  
**Active Fibronectin Type III Domain Containing Protein 5 (FNDC5)**  
**Organism Species: Mus musculus (Mouse)**  
***Instruction manual***

FOR RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

13th Edition (Revised in Aug, 2023)

## **[ PROPERTIES ]**

**Source:** Eukaryotic expression.

**Host:** 293F cell

**Residues:** Asp29~Glu140

**Tags:** N-terminal His-tag

**Purity:** >90%

**Endotoxin Level:** <1.0EU per 1µg (determined by the LAL method).

**Buffer Formulation:** PBS, pH7.4, containing 5% Trehalose.

**Applications:** Cell culture; Activity Assays; In vivo assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 4.7

**Predicted Molecular Mass:** 14.2kDa

**Accurate Molecular Mass:** 22-30kDa as determined by SDS-PAGE reducing conditions.

### **Phenomenon explanation:**

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## **[ USAGE ]**

Reconstitute in ddH<sub>2</sub>O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


## **[ SEQUENCE ]**

```
DS PSAPVNVTVR HLKANSVVV  
WDVLEDEVVI GFAISQQKKD VRMLRFIQEV NTTTRSCALW DLEEDTEYIV  
HVQAISIQQG SPASEPVLFK TPREAEMAS KNKDEVTMKE
```

## **[ ACTIVITY ]**

Fibronectin type III domain-containing protein 5, the precursor of irisin, is a protein that is encoded by the FNDC5 gene. FNDC5 is a 12 kDa glycosylated polypeptide hormone that regulate glycolipid metabolism, cardiovascular homeostasis metabolism, stem cell differentiation, and neuronal development. Brain Derived Neurotrophic Factor (BDNF) has been identified as an interactor of FNDC5, thus a functional binding ELISA assay was conducted to detect the interaction of recombinant mouse FNDC5 and recombinant dog BDNF. Briefly, FNDC5 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100  $\mu$ l were then transferred to BDNF-coated microtiter wells and incubated for 1h at 37 °C . Wells were washed with PBST and incubated for 1h with anti-FNDC5 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37 °C , wells were aspirated and washed 5





**Figure 3. SDS-PAGE**

**Sample: Active recombinant FNDC5, Mouse**

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.