



**APA062Hu01 100µg**

**Active Interleukin 16 (IL16)**

**Organism Species: Homo sapiens (Human)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Met1203~Ser1332

**Tags:** N-terminal His-tag

**Purity:** >98%

**Buffer Formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

**Applications:** Cell culture; Activity Assays; In vivo assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 6.0

**Predicted Molecular Mass:** 17.1kDa

**Accurate Molecular Mass:** 17kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**


```
MPDLNSST DSAASASAAS DVSVESTAEA TVCTVTLEKM SAGLGFSLEG  
GKGSLSHGDKP LTINRIFKGA ASEQSETVQP GDEILQLGGT AMQGLTRFEA  
WNIIKALPDG PVTIVIRRKS LQSKETTAAG DS
```

## **[ ACTIVITY ]**

Pro-IL16 (Interleukin16) is a 631 amino acid precursor molecule, which is then cleaved into different isoforms. Researches have shown that the recombinant human IL16, containing C-terminal 130 amino acids, has same bioactivity as the natural secreted human IL16. Besides, IL16 has been considered to stimulate the proliferation of Jurkat cells at low dose ( $10^{-9}$  M). Thus, a proliferation assay of recombinant human IL16 was conducted using Jurkat cells. Briefly, Jurkat cells were seeded into triplicate wells of 96-well plates at a density of 10, 000 cells/well in RPMI-1640 with the addition of various concentrations of IL16. After incubated for 72h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 $\mu$ L of CCK-8 solution was added to each well of the plate, then the absorbance at 450nm was measured using a microplate reader after incubating the plate for 1-4 hours at 37°C. Cell proliferation of Jurkat cells after incubation with IL16 for 72h observed by inverted microscope was shown in Figure 1. The CCK-8 data was shown in Figure 2. It was obvious that IL16 significantly promoted cell proliferation of Jurkat cells.




**Figure 1. Cell proliferation of Jurkat cells after stimulated with IL16.**




**Figure 2. Cell proliferation of Jurkat cells after stimulated with IL16.**

## [ IDENTIFICATION ]



**Figure 3. SDS-PAGE**

**Sample: Active recombinant IL16, Human**



**Figure 4. Western Blot**

**Sample: Recombinant IL16, Human;**

**Antibody: Rabbit Anti-Human IL16 Ab (PAA062Hu01)**

### **[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.