

**APB837Mu02 100µg**  
**Active Interleukin 2 Receptor Alpha (IL2Ra)**  
**Organism Species: Mus musculus (Mouse)**  
*Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

1st Edition (Apr, 2016)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Glu22~Lys236

**Tags:** N-terminal His-tag

**Purity:** >92%

**Buffer Formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

**Applications:** Cell culture; Activity Assays; In vivo assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 7.2

**Predicted Molecular Mass:** 28.4kDa

**Accurate Molecular Mass:** 37kDa as determined by SDS-PAGE reducing conditions.

### **Phenomenon explanation:**

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

## [ USAGE ]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ SEQUENCE ]

```
ELCLYDPPE VPNATFKALS YKNGTILNCE
CKRGFRRLKE LVYMRCLGNS WSSNCQCTSN SHDKSRKQVT AQLEHQKEQQ
TTTDMQKPTQ SMHQENLTGH CREPPPWKHE DSKRIYHFVE GQSVHYECIP
GYKALQRGPA ISICKMKCGK TGWTQPQLTC VDEREHRFL ASEESQGSRN
SSPESETSCP ITTTDFPQPT ETTAMTETFV LTMEYK
```

## [ ACTIVITY ]

IL2RA (Interleukin-2 receptor alpha chain ), also known as Tac antigen and as CD25, was initially identified as a 55kDa membrane glycoprotein which is capable of binding IL-2. Besides, mouse IL2 shares 75.1% AA sequence identity with rat IL2, suggesting the exist of cross-species activity. Thus we have conducted a binding ELISA assay to detect the interaction of recombinant rat IL2RA with recombinant mouse IL2. Briefly, IL2RA were diluted serially in PBS, with 0.01%BSA (pH 7.4). Duplicate samples of 100uL were then transferred to IL2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-IL2RA pAb, then aspirated and washed 3

times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of IL2RA with IL2 was shown in Figure 1 and this effect was in a dose dependent manner.




Figure 1. The binding activity of IL2RA with IL2.

## [ IDENTIFICATION ]





Figure 2. SDS-PAGE

Sample: Active recombinant IL2Ra, Mouse



**Figure 3. Western Blot**

**Sample: Recombinant IL2Ra, Mouse;**

**Antibody: Rabbit Anti-Mouse IL2Ra Ab (PAB837Mu02)**