

**APA553Mu61 50µg**  
**Active Matrix Metalloproteinase 9 (MMP9)**  
**Organism Species: *Mus musculus (Mouse)***  
***Instruction manual***

FOR RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

13th Edition (Revised in Aug, 2023)

## **[ PROPERTIES ]**

**Source:** Eukaryotic expression.

**Host:** 293F cell

**Residues:** Ser225~Asp390

**Tags:** N-terminal His-tag

**Purity:** >90%

**Endotoxin Level:** <1.0EU per 1µg (determined by the LAL method).

**Buffer Formulation:** PBS, pH7.4, containing 5% Trehalose.

**Original Concentration:** 50µg/mL

**Applications:** Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 5.7

**Predicted Molecular Mass:** 19.9kDa

**Accurate Molecular Mass:** 19kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-0.2 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the

protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**

```
SNGAPC HFPFTFEGRS YSACTTDGRN  
DGTPWCSTTA DYDKDGKFGF CPSELYTEH GNGEGKPCVF PFIFEGRSYS  
ACTTKGRSDG YRWCATTANY DQDKLYGFCP TRVDATVVGG NSAGELCVFP  
FVFLGKQYSS CTSDGRRDGR LWCATTSNFD TDKKWGFCPD
```

## **[ ACTIVITY ]**

Mechanism: MMP9 is a zinc-dependent enzymes capable of cleaving components of the extracellular matrix, which belongs to the matrix metalloproteinase (MMP) family . It is a gelatinase A, 92 kDa type IV collagenase which can hydrolyze gelatin under certain conditions. Gelatin zymography is mainly used for the detection of the gelatinases, MMP-2 and MMP-9 and It is extremely sensitive because levels of 10 pg of MMP-2 can already be detected .Briefly , various concentrations of recombinant mouse MMP9 (100ng, 50ng, 25ng, 12.5ng, 6.25ng, 3.1ng, 1.5ng and 0.7ng) were denatured by SDS loading buffer, electrophoresed through sodium dodecylsulphate - polyacrylamide gel (SDS - PAGE; 15% gels) containing gelatin (1 mg/ml) with nonreducing conditions. After renaturation, incubation and CCB-stained, active MMP9 would hydrolyze gelatin nearby, which was indicated by the white binds on the gel. In this experiment we use trypsin as positive control.


Result: Gelatin hydrolysis by recombinant mouse MMP9(10-70kd) was shown in figure 1.

Marker 100 50 25 12.5 6.25 3.1 1.5 0.7ng positive



**Figure 1**

## **[ IDENTIFICATION ]**



**Figure 2. SDS-PAGE**

**Sample: Active recombinant Mouse, MMP9**

## **[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.