

APA679Hu01 100µg

Active Protein Kinase, AMP Activated Alpha 1 (AMPK Alpha 1)

Organism Species: *Homo sapiens* (Human)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Ile28~Lys257

Tags: N-terminal His-tag

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 200µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 8.5

Predicted Molecular Mass: 29.6kDa

Accurate Molecular Mass: 31kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[**SEQUENCE**]

```
ILG DTLGVGTFGK VKVGKHELTG
HKVAVKILNR QKIRSLDVVG KIRREIQNLK LFRHPHIKL YQVISTPSDI
FMVMEYVSGG ELFDYICKNG RLDEKESRRL FQQILSGVDY CHRHMVVHRD
LKPENVLLDA HMNAKIADFG LSNMMSDGEF LRTSCGSPNY AAPEVISGRL
YAGPEVDIWS SGVILYALLC GTLPFDDDHV PTLFKKICDG IFYTPQYLNPN
SVISLLK
```

[**ACTIVITY**]

AMPK alpha 1, also known as AMP-activated protein kinase alpha 1 subunit, is a crucial enzyme that plays a significant role in energy metabolism. It is involved in regulating several metabolic processes, including glucose and fatty acid uptake, oxidation, and storage. It promotes energy-producing pathways, such as glycolysis, fatty acid oxidation, and mitochondrial biogenesis, while inhibiting energy-consuming processes, like protein synthesis and lipogenesis. Besides, PRKAb1 has been identified as an interactor of AMPK Alpha 1, thus a functional binding ELISA assay was conducted to detect the interaction of recombinant human AMPK Alpha 1 and recombinant human PRKAb1. Briefly, AMPK Alpha 1 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to PRKAb1-coated microtiter wells and incubated for 1h at 37°C. Wells were washed with PBST and incubated for 1h with anti-AMPK Alpha 1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37°C, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 μ l stop solution to the wells and read at 450/630 nm immediately.

The binding activity of recombinant human AMPK Alpha 1 and recombinant human PRKAb1 was shown in Figure 1, the EC50 for this effect is 0.94 ug/mL.




Figure 1. The binding activity of recombinant human AMPK Alpha 1 and recombinant human PRKAb1

[IDENTIFICATION]

ATTCGGGTGACAGCCTGGGGTGGCCCTTGGGAAAGTGAAGGTTGGAAACATGAAATGACTG99CATAAAGTACTGTGAAAGTACTCAATGGACAGAAATGGAGCCTTGAATGTGTAAGGAAATTCAGAACTCAGCTTTTCAGGCATCTCATATAATTAA
 I L G D T L G V G T F G K V K V G K H E L T G H K V A V K I L N R Q K I R S L D V V G K I R R E I Q N L K L F R H P H I I K




Figure 2. Gene Sequencing (extract)




Figure 3. SDS-PAGE

Sample: Active recombinant AMPK Alpha 1, Human

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.