

**APA567Mu01 50µg**

**Active S100 Calcium Binding Protein B (S100B)**

**Organism Species: Mus musculus (Mouse)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

13th Edition (Revised in Aug, 2023)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Met1~Glu92

**Tags:** N-terminal His-tag

**Purity:** >92%

**Endotoxin Level:** <1.0EU per 1µg (determined by the LAL method).

**Buffer Formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 0.01% skl, 5%Trehalose.

**Applications:** Cell culture; Activity Assays; In vivo assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 4.9

**Predicted Molecular Mass:** 12.0kDa

**Accurate Molecular Mass:** 10kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**

```
MSELEKAMVA LIDVFHQYSG REGDKHKLKK SELKELINNE LSHFLEEIKE  
QEVVDKVMET LDEDGDGECDFQEFMAFVAM VTTACHEFFE HE
```

## **[ ACTIVITY ]**

Protein S100B is a member of the S100 family. S100 proteins are EF-hand calcium-binding proteins and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. Experimental results suggest that the receptor for advanced glycation end products (RAGE) plays important roles in mediating S100 protein-induced cellular signaling. Besides, mouse RAGE shares similarities with human RAGE in amino acids sequence with the identity of 78.2%. Thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse S100B and recombinant human RAGE. Briefly, S100B were diluted serially in PBS, with 0.01%BSA (pH 7.4). Duplicate samples of 100uL were then transferred to RAGE-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-S100B mAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50μL stop solution to the wells and read at 450nm immediately. The binding activity of S100B and RAGE was shown in Figure 1, and this effect was in a dose dependent manner.




Figure 1. The binding activity of S100B with RAGE.

## [ IDENTIFICATION ]




Figure 2. SDS-PAGE

Sample: Active recombinant S100B, Mouse

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.