

**APA552Ov01 100µg**  
**Active Tissue Inhibitors Of Metalloproteinase 1 (TIMP1)**  
**Organism Species: *Ovis aries*; *Ovine* (Sheep)**  
***Instruction manual***

FOR RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

13th Edition (Revised in Aug, 2023)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Cys24~Cys197

**Tags:** N-terminal His-tag

**Purity:** >90%

**Endotoxin Level:** <1.0EU per 1µg (determined by the LAL method).

**Buffer Formulation:** PBS, pH7.4, containing 0.01% SKL, 5%Trehalose .

**Original Concentration:** 200µg/mL

**Applications:** Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 8.0

**Predicted Molecular Mass:** 23.2kDa

**Accurate Molecular Mass:** 23kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**

```
CTCVPPH PQTAFCNSEV VIRAKFVGTA EVNETALYQR YEIKMTKMPK  
GFSALRDAPD IRFIYTPAME SVCGYFHRSQ NRSEEFLIAG QLSNGHLHIT  
TCSFVAPWNS MSSAQRRGFT KTYAAGCEEC TVFPCSSIPK KLQSDTHCLW  
TDQLLTGSDK GFQSRHLACL PREPGMC
```

## **[ ACTIVITY ]**

Tissue inhibitors of metalloproteinase 1 (TIMP1) is a member of the family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). TIMP-1 is a glycoprotein with a molecular mass of 28 kDa produced by a wide range of cell types. TIMP-1 inhibits active MMP-mediated proteolysis by forming an N-terminal, non-covalent binary complex with the MMP active site. The activity of recombinant sheep TIMP1 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH<sub>2</sub> in the assay buffer 50 mM Tris, 10 mM CaCl<sub>2</sub>, 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM APMA at 37 ° C for 1 hour and roTIMP1 (MW: 23.2 KD) was diluted to different concentrations with the assay buffer. Mix 8 µl of roTIMP1 curve dilutions, 12.8 µl of activated rhMMP-2, and 59.2 µl of assay buffer, including a control containing assay buffer and the diluted rhMMP-2 and incubate the reactions for 2 hours at 37 ° C. Loading 50 µl of the incubated mixtures which were diluted five-fold in assay buffer into empty wells of a plate, and start the reaction by adding 50 µl of 20 µM substrate. Include a substrate blank containing 50 µl of assay buffer and 50 µl of 20 µM substrate. Then read at excitation and

emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown in Figure 1 and it was obvious that recombinant sheep TIMP1 significantly decreased rhMMP2 activity. The inhibition IC50 was <23 nM.




Figure 1. Inhibition of MMP2 activity by recombinant sheep TIMP1

## [ IDENTIFICATION ]




Figure 2. SDS-PAGE

**Sample: Active recombinant TIMP1, Sheep**

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.