

APA128Ra01 100µg
Active Tissue Inhibitors Of Metalloproteinase 2 (TIMP2)
Organism Species: *Rattus norvegicus* (Rat)
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Cys27~Pro220

Tags: N-terminal His-tag

Purity: >90%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5%Trehalose .

Original Concentration: 200µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.5

Predicted Molecular Mass: 23.0kDa

Accurate Molecular Mass: 23kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
CSCS PVHPQQAFCN ADVVIRAKAV  
SEKEVDSGND IYGNPIKRIQ YEIKQIKMFK GPKDIEFIY TAPSSAVCGV  
SLDVGGKKEY LIAGKAEGDG KMHITLCDFI VPWDTLSITQ KKSLSNHRYQM  
GCECKITRCP MIPCYISSPD ECLWMDWVTE KSINGHQAKF FACIKRSDGS  
CAWYRGAAPP KQEFLDIEDP
```

[ACTIVITY]

Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs). There are four members of the family, TIMP-1, TIMP-2, TIMP-3, and TIMP-4. TIMP-2 is a non N-glycosylated protein with a molecular mass of 22 kDa produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes. TIMP-2 also has erythroid-potentiating and cell growth promoting activities. The activity of recombinant rat TIMP2 was measured by its ability to inhibit rhMMP2 cleavage of a fluorogenic peptide substrate MCA-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH₂ in the assay buffer 50 mM Tris, 10 mM CaCl₂, 150 mM NaCl, 0.05% (w/v) Brij-35, pH 7.5. rhMMP2 was diluted to 100 ug/ml and activated with 1 mM APMA at 37 ° C for 1 hour and rrTIMP2 (MW: 22.98 KD) was diluted to different concentrations with the assay buffer. Mix 8 µl of rrTIMP2 curve dilutions, 12.8 µl of activated rhMMP-2, and 59.2 µl of assay buffer, including a control containing assay buffer and the diluted rhMMP-2 and incubate the reactions for 2 hours at 37 ° C. Loading 50 µl of the incubated mixtures which were diluted five-fold in assay buffer into empty wells of

a plate, and start the reaction by adding 50 μ l of 20 μ M substrate. Include a substrate blank containing 50 μ l of assay buffer and 50 μ l of 20 μ M substrate. Then read at excitation and emission wavelengths of 320 nm and 405 nm, respectively, in kinetic mode for 5 minutes. The result was shown in Figure 1 and it was obvious that recombinant rat TIMP2 significantly decreased rhMMP2 activity. The inhibition IC₅₀ was <1.7 nM.




Figure 1. Inhibition of MMP2 activity by recombinant rat TIMP2

[IDENTIFICATION]




Figure 2. SDS-PAGE

Sample: Active recombinant TIMP2, Rat

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.