



A90852Mu22

Antibody to Protease Activated Receptor 2 (PAR2)

Organism: Mus musculus (Mouse)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

6th Edition (Revised in March, 2013)

[PRODUCT INFORMATION]

Immunogen: PAR2, Mouse

Clonality: Monoclonal

Host: Rat


Immunoglobulin Type: IgG

Purification: Affinity Chromatography. **Applications:** WB, ICC, IHC-P, IHC-F

, ,

Concentration: 500µg/mL

UOM: 100μg



Sample: Recombinant PAR2, Mouse

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant PAR2 (Met214~Tyr325) expressed in *E.coli*.

USCN Accession No.: P90852Mu01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence

is listed below.

MGHHHHHHSGSEF- MKQTIYI PALNITTCHD VLPEEVLVGD MFNYFLSLAI GVFLFPALLT ASAYVLMIKT LRSSAMDEHS EKKRQRAIRL IITVLAMYFI CFAPSNLLLV VHYFLIKTQR OSHVY





[ANTIBODY SPECIFITY]

Anti PAR2 is a rat monoclonal antibody raised against PAR2. It has been selected for its ability to recognize PAR2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant PAR2 (Met214~Tyr325) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.