

PAC038Ge71

Biotin-linked Antibody to Diacylglycerol (DAG)

Organism Species: General

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: DAG-BSA

Clonality: Polyclonal

Conjugation: Biotin

Host: Rabbit

Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL


UOM: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Small Molecule, DAG conjugated to BSA.

Accession No.: CPC038Ge11

Structural Formula:



Chemical structure of the diglyceride 1-palmitoyl-2-oleoyl-glycerol

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against DAG. It has been selected for its ability to recognize DAG in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN_3 , 50% glycerol.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.