

MAJ085Hu21

Monoclonal Antibody to Follistatin Like Protein 1 (FSTL1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C7

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 100µl

**Cross Reactivity: Porcine** 

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant FSTL1 (Glu21~lle308) expressed in E.coli

Accession No.: RPJ085Hu01

#### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

glycerol.

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50%

# [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

# Coud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]

kDa	
70-	
44-	
33- 27- 22-	
18-	
14– 10–	

lysate; Primary Ab: 2µg/ml Mouse Anti-Human FSTL1 Antibody Second Ab: 0.2µg/ml HRP-Linked Caprine Anti-

Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

DAB staining on IHC-P; Samples:

Western Blot; Sample: Human Placenta Human Cerebrum Tissue; Primary Ab: 10µg/ml Mouse Anti-Human FSTL1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.