

**MAA421Hu21** 

Monoclonal Antibody to Myelin Oligodendrocyte Glycoprotein (MOG)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C2

Traits: Liquid

Concentration: 1mg/mL

**UOM:** 100µL

Cross Reactivity: Rat; Porcine.

Applications: WB; IHC

### [ IMMUNOGEN ]

Immunogen: Recombinant MOG (Gly30~Tyr149) expressed in E.coli

Accession No.: RPA421Hu01

#### [ APPLICATIONS ]

Western blottin:g: 0.01-2µg/mL;

Immunohistochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



Western Blot; Sample: Lane1: Rat
Cerebrum lysate; Lane2: Rat
Cerebellum lysate Primary Ab: 2µg/ml
Mouse Anti-Human MOG Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody (Catalog: SAA544Mu19)



DAB staining on IHC-P; Sample:
Human Cerebrum Tissue; Primary Ab:
30µg/ml Mouse Anti-Human MOG
Antibody Second Ab: 2µg/mL HRPLinked Caprine Anti-Mouse IgG
Polyclonal Antibody (Catalog:
SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.