

PAB054Mu01

Polyclonal Antibody to Apoptotic Peptidase Activating Factor 1 (APAF1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µl

Cross Reactivity: Rat

Applications: WB; IHC; IF

#### [ IMMUNOGEN ]

Immunogen: Recombinant APAF1 (Arg872~Phe1135 (Accession # O88879)) expressed in

E.coli

Accession No.: RPB054Mu01

#### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

## Coud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]





Figure. Western Blot; Sample: Recombinant APAF1, Mouse.




Mouse Cardiac Muscle Tissue; Primary Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:

SAA544Rb19)


DAB staining on IHC-P; Samples:

Samples: Mouse Kidney Tissue; Ab: 20ug/ml Rabbit Anti-Mouse APAF1 Primary Ab: 20ug/ml Rabbit Anti-Mouse APAF1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

DAB staining on IHC-P;




DAB staining on IHC-P; Sample: Mouse Kidney Tissue



DAB staining on IHC-P;

Sample: Mouse Cardiac Muscle Tissue



Samples: Human Hela cell;

FITC staining on IF;

APAF1 Antibody Control: Used PBS instead of primary antibody

Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody


Primary Ab: 20µg/ml Rabbit Anti-Mouse Primary Ab: 20µg/ml Rabbit Anti-Mouse **APAF1** Antibody Control: Used PBS instead of primary antibody Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

Primary Ab: 20µg/ml Rabbit Anti-Mouse APAF1 Antibody Second Ab: 1.5µg/ml FITC-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb18)

# Coud-Clone Corp.

(Catalog: SAA544Rb19)

(Catalog: SAA544Rb19)



FITC staining on IF;

Sample: Hela cell

Primary Ab: 20µg/ml Rabbit Anti-Mouse

**APAF1** Antibody

Second Ab: 2µg/ml FITC-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb11)

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.