

PAA040Ra01

Polyclonal Antibody to Chemokine C-X3-C-Motif Ligand 1 (CX3CL1)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/ml

UOM: 50µl

Cross Reactivity: Human

Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant CX3CL1 (Gln83~Pro328) expressed in E.coli

Accession No.: RPA040Ra01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunofluorescence:5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]







Figure. Western Blot; Sample: Recombinant CX3CL1, Rat.




DAB staining on IHC-P; Sample: Rat Small intestine Tissue; Primary Ab: 20ug/ml Rabbit Anti-Rat CX3CL1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)






DAB staining on IHC-P;
Sample: Rat Spleen Tissue;
Primary Ab: 20ug/ml Rabbit Anti-Rat
CX3CL1 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Rat Small intestine Tissue
Primary Ab: 20µg/ml Rabbit Anti-Rat
CX3CL1 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody





DAB staining on IHC-P;
Sample: Rat Lung Tissue
Primary Ab: 20µg/ml Rabbit Anti-Rat
CX3CL1 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody


(Catalog: SAA544Rb19)


(Catalog: SAA544Rb19)

్రి Cloud-Clone Corp.











DAB staining on IHC-P; Sample: Rat Spleen Tissue Primary Ab: 20µg/ml Rabbit Anti-Rat CX3CL1 Antibody Control: Used PBS instead of primary antibody Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal

> Antibody (Catalog: SAA544Rb19)

Western Blot; Sample: Rat Heart lysate; Western Blot; Sample: Rat Cerebrum Primary Ab: 1µg/ml Rabbit Anti-Rat CX3CL1 Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

lysate; Primary Ab: 1µg/ml Rabbit Anti-Rat CX3CL1 Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)





Western Blot; Sample: Rat Kidney lysate:

Primary Ab: 1µg/ml Rabbit Anti-Rat CX3CL1 Antibody

Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)

Western Blot; Sample: Human PC3 cell

lysate:

Primary Ab: 1µg/ml Rabbit Anti-Rat

CX3CL1 Antibody

Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)

[IMPORTANT NOTE]



The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.