

PAB931Hu01

**Polyclonal Antibody to Fumarase (FUM)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM**: 100µl

**Cross Reactivity: Porcine** 

Applications: WB,IHC,ICC/IF

#### [ IMMUNOGEN ]

Immunogen: Recombinant FUM (Ala45~Asn188) expressed in E.coli

Accession No.: RPB931Hu01

### [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL

Immunohistochemistry: 5-20µg/mL

Immunofluorescence:5-20µg/mL

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]







Figure. Western Blot; Sample: Recombinant FUM, Human.




DAB staining on IHC-P; Samples:
Human Liver Tissue; Primary Ab:
20µg/ml Rabbit Anti-Human FUM
Antibody Second Ab: 2µg/mL HRPLinked Caprine Anti-Rabbit IgG
Polyclonal Antibody (Catalog:
SAA544Rb19)




DAB staining on IHC-P;
Samples: Human Glioma Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Samples: Human Kidney Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Samples: Human Stomach Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;
Sample: Human Liver Tissue
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

# Cloud-Clone Corp.



DAB staining on IHC-P;
Sample: Human Kidney Tissue
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody



(Catalog: SAA544Rb19)

**Knockout Varification:** 

Lane 1: Wild-type Hela cell lysate;

Lane 2: FUM knockout Hela cell lysate;

Predicted MW: 55,50kd

Observed MW: 50kd


Primary Ab: 1µg/ml Rabbit Anti-Human

**FUM Antibody** 

Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal


Antibody



DAB staining on IHC-P;
Sample: Human Stomach Tissue
Primary Ab: 20µg/ml Rabbit AntiHuman FUM Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked

Antibody
(Catalog: SAA544Rb19)

Caprine Anti-Rabbit IgG Polyclonal



Samples: Human Hela Cells; Primary Ab: 20µg/ml Rabbit Anti-Human FUM Antibody Second Ab: 1µg/ml FITC-Linked


FITC staining on IF;

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb18)

(Catalog: SAA544Rb19)



Western Blot; Sample: Lane1: Hela cell lysate; Lane2: Porcine Liver lysate Primary Ab: 1µg/ml Rabbit Anti-Human FUM Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)



(Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.