

PAB547Ra01

Polyclonal Antibody to Indoleamine-2,3-Dioxygenase (IDO)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

### [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µL

Cross Reactivity: Human; Mouse

Applications: WB; IHC; ICC; IP.

#### [<u>IMMUNOGEN</u>]

Immunogen: Recombinant IDO (Ser276~Lys360) expressed in E.coli

Accession No.: RPB547Ra01

#### [APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [IDENTIFICATION]

kDa 100–	kDa 100-
70-	70-
44-	44-
33– 27–	33- 27-
22-	22-
Western Blot; Sample: Rat Placenta	Western Blot; Sample: Human Placenta
lysate;	lysate;
Primary Ab: 2µg/ml Rabbit Anti-Rat IDOPrimary Ab: 2µg/ml Rabbit Anti-Rat IDO	
Antibody	Antibody
Second Ab: 0.2µg/mL HRP-Linked	Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal	Caprine Anti-Rabbit IgG Polyclonal
Antibody	Antibody
(Catalog: SAA544Rb19)	(Catalog: SAA544Rb19)

### [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.