

PAC614Ra01

Polyclonal Antibody to Macrophage Receptor With Collagenous Structure (MARCO)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Cond-Clone Corp.

## [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µL

Cross Reactivity: Mouse; Porcine

Applications: WB; IHC; ICC; IP.

#### [<u>IMMUNOGEN</u>]

Immunogen: Recombinant MARCO (Gly367~Arg519) expressed in E.coli

Accession No.: RPC614Ra01

#### [APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no


## Cloud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


## [IDENTIFICATION]









Vector Red staining on IHC-P; Samples: Human Thymus Tissue; Primary Ab: 20µg/ml Rabbit Anti-Human MARCO Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



Western Blot; Sample: Human Leukocyte lysate; Primary Ab: 1µg/ml Rabbit Anti-Rat MARCO Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19) Selected





Western Blot; Samples: Lane1: Porcine	Western Blot; Sample: RAW264.7 cell
Lung lysate; Lane2: Porcine Lymph	lysate
node lysate;	Primary Ab: 0.5µg/ml Rabbit Anti-Rat
Primary Ab: 0.5µg/ml Rabbit Anti-Rat	MARCO Antibody
MARCO Antibody	Second Ab: 0.2?g/ml HRP-Linked
Second Ab: 0.2?g/ml HRP-Linked	Caprine Anti-Rabbit IgG Polyclonal
Caprine Anti-Rabbit IgG Polyclonal	Antibody
Antibody	(Catalog: SAA544Rb19)
(Catalog: SAA544Rb19)	



### [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.