

PAA553Mu01

Polyclonal Antibody to Matrix Metalloproteinase 9 (MMP9)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [ PROPERTIES ]

**Source:** Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 500µg/mL

**UOM:** 100µL

Cross Reactivity: Human

**Applications:** WB,ICC/IF

#### [ IMMUNOGEN ]

Immunogen: Recombinant MMP9 (Ser225~Asp390) expressed in E.coli

Accession No.: RPA553Mu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunofluorescence:5-20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

# Coud-Clone Corp.

expiration date under appropriate storage condition.

#### [ IDENTIFICATION ]





Figure. Western Blot; Sample: Recombinant MMP9, Mouse.



Western Blot; Sample: Human Jurkat cell lysate;

Primary Ab: 1µg/mL Rabbit Anti-Mouse

MMP9 Antibody

Second Ab: 0.2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)



FITC staining on IF;
Samples: Human HepG2 cell;
Primary Ab: 20µg/mL Rabbit AntiMouse MMP9 Antibody
Second Ab: 1.5µg/mL FITC-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb18)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.