

PAA886Rb51

Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma (PPAR γ)

Organism Species: *Oryctolagus cuniculus* (Rabbit)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Guinea pig

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Traits: Liquid

Concentration: 0.5mg/ml

UOM: 200µl

Cross Reactivity: Mouse

Applications: WB; IHC; ICC; IP.

[IMMUNOGEN]

Immunogen: Recombinant PPARg (Phe280~Tyr475 (Accession # O19052)) expressed in *E.coli*

Accession No.: RPA886Rb01

[APPLICATIONS]

Western blotting: 0.5-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunocytochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]





Figure. Western Bolt; Sample: Recombinant PPARγ, Rabbit.




Western Bolt; Sample: Mouse Liver lysate

Primary Ab: 1?g/ml Cavia Anti-Rabbit PPARγ Antibody

Second Ab: 0.2μg/mL HRP-Linked Rabbit Anti-Cavia IgG Polyclonal Antibody

(Catalog: SAA544Gu09)



DAB staining on IHC-P;

Samples: Rabbit Cardiac Muscle Tissue;

Primary Ab: 10?g/ml Cavia Anti-Rabbit PPARγ Antibody

Second Ab: 2μg/mL HRP-Linked Rabbit Anti-Cavia IgG Polyclonal Antibody (Catalog: SAA544Gu09)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.