

**PAL096Hu01** 

Polyclonal Antibody to Programmed Cell Death Protein 6 (PDCD6)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µl

**Cross Reactivity:** 

Applications: WB; IHC; ICC/IF; FCM

[ IMMUNOGEN ]

Immunogen: Recombinant PDCD6 (Phe32~Val191) expressed in E.coli

Accession No.: RPL096Hu01

[ APPLICATIONS ]

Western blotting: 0.01-2µg/mL

Immunohistochemistry: 5-20µg/mL

Immunocytochemistry: 5-20µg/mL

Flow cytometry:20µg/ml;

Optimal working dilutions must be determined by end user.

### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

# Coud-Clone Corp.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]







Figure. Western Blot; Sample: Recombinant PDCD6, Human.



FITC staining on IF; Samples: Human Hela cell; Primary Ab: 20µg/ml Rabbit Anti-Human PDCD6 Antibody Second Ab: 1µg/ml FITC-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb18)




DAB staining on IHC-P;
Samples: Human Kidney Tissue;
Primary Ab: 20µg/ml Rabbit AntiHuman PDCD6 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Hela cell lysate
Primary Ab: 1µg/ml Rabbit Anti-Human
PDCD6 Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)




Knockout Varification:

Lane 1: Wild-type Hela cell lysate;

Lane 2: PDCD6 knockout Hela cell lysate;

Predicted MW: 23.7kd

Predicted MW: 22,7kd Anti-human PDCD6 Polyclonal
Observed MW: 23kd Antibody (Catalog PAL096Hu01, red
Primary Ab: 1µg/ml Rabbit Anti-Human histogram) or Isotype control antibody
PDCD6 Antibody (Catalog IS067-Rb01, green



Human Hela cell was fixed with 2% paraformaldehyde (10 min), permeabilised with 0.1% BSA-Triton X-100,then stained with 20µg/ml rabbit Anti-human PDCD6 Polyclonal Antibody (Catalog PAL096Hu01, red histogram) or Isotype control antibody (Catalog IS067-Rb01, green



Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

histogram), followed by 1µg/ml FITC-conjugated Anti-rabbit IgG Secondary
Antibody (Catalog SAA544Rb18).

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.