

PAB537Hu01

Polyclonal Antibody to Protease, Serine 2 (PRSS2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



### [PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 100µL

Cross Reactivity: Mouse; Rat; Porcine

Applications: WB

#### [ IMMUNOGEN ]

Immunogen: Recombinant PRSS2 (Ile24~Asp241) expressed in E.coli

Accession No.: RPB537Hu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



## [ IDENTIFICATION ]






Figure. Western Blot; Sample: Recombinant PRSS2, Human.



Western Blot; Sample: Porcine
Pancreas lysate
Primary Ab: 0.1µg/ml Rabbit AntiHuman PRSS2 Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Lane1: Rat
Pancreas lysate; Lane2: Mouse
Pancreas lysate
Primary Ab: 0.1µg/ml Rabbit AntiHuman PRSS2 Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.