

PA	NΩ	15	۸a	Λı	ı∩1
			O I I	,,,,	

Polyclonal Antibody to T-Cell Immunoreceptor With Ig And ITIM Domains Protein (TIGIT)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



# [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.48mg/mL

**UOM:** 50µL

Cross Reactivity: N/A

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant TIGIT (Gly26~Ser238) expressed in E.coli

Accession No.: RPN056Mu01

### [ APPLICATIONS ]

Western blotting: 0.01-3µg/mL

Immunohistochemistry: 5-20µg/mL

Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no



obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [ IDENTIFICATION ]





DAB staining on IHC-P; Sample: Mouse Western Blot; Sample: Mouse Spleen

Spleen Tissue; Primary Ab: 20ug/ml

Rabbit Anti-Mouse TIGIT Antibody

Second Ab: 2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody (Catalog: SAA544Rb19)

lysate

Primary Ab: 3µg/ml Rabbit Anti-Mouse

**TIGIT Antibody** 

Second Ab: 0.2?g/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.